

c-Met antibody

Cat. No. GTX10728

Host	Goat	
Clonality	Polyclonal	
Isotype	IgG	
Applications	WB, FCM, IHC, Neutralizing/Inhibition	
Reactivity	Human	

Package 25 μg

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.2 μg/ml
FCM	Assay dependent
IHC	5-15 μg/ml
Neutralizing/Inhibition	0.5-2 μg/ml

Not tested in other applications.

Calculated MW 156 kDa. (Note)

Properties		
Form	Liquid	
Buffer	PBS carbohydrates	
Preservative	No preservatives	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)	
Immunogen	purified recombinant human hepatocyte growth factor receptor extracellular domain expressed in mouse NSO cells.	
Purification	Purified by affinity chromatography	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 1 of 2



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 2 of 2