

## c-Met antibody

**Cat. No. GTX10728**

<b>Host</b>	Goat
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, FCM, IHC, Neutralizing/Inhibition
<b>Reactivity</b>	Human

**Package**

25 µg

### Applications

#### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	0.1-0.2 µg/ml
FCM	Assay dependent
IHC	5-15 µg/ml
Neutralizing/Inhibition	0.5-2 µg/ml

Not tested in other applications.

**Calculated MW** 156 kDa. ([Note](#))

### Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS carbohydrates
<b>Preservative</b>	No preservative
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	Batch dependent (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	purified recombinant human hepatocyte growth factor receptor extracellular domain expressed in mouse NSO cells.
<b>Purification</b>	Purified by affinity chromatography
<b>Conjugation</b>	Unconjugated

#### Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).



For full product information, images and publications, please visit our [website](#).