

BAK antibody [TC100]

Cat. No. GTX10808

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Application	WB, IHC-Fr
Reactivity	Human

Package

50 µg

APPLICATION

Application Note

IHC-Fr: Use at a concentration of 1 - 5 µg/ml. WB: Use at a concentration of 2 - 5 µg/ml. Detects a band of approximately 30 kDa (predicted molecular weight: 23.4 kDa). Optimal dilutions/concentrations should be determined by the end user.

Calculated MW 23 kDa. ([Note](#))


PROPERTIES

Form	Liquid
Buffer	50mM Sodium Phosphate, 0.2% Gelatin
Preservative	0.09% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment (C-terminal transmembrane portion deleted) (Human).
Purification	Immunogen affinity purified
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption. Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX10808 WB Image

Detection of human Bak by immunoblotting. Sample: Whole cell lysates from untreated HL-60 cells. Primary antibody: Anti-Bak (Ab-1) Mouse mAb (TC-100) (GTX10808) (2 µg/ml). Detection: Chemiluminescence.



For full product information, images and publications, please visit our [website](https://www.genetex.com).