

# Histone H2A.X antibody

## Cat. No. GTX108272

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human, Mouse, Rat



## Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:20000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500

Not tested in other applications.

**Calculated MW** 15 kDa. ( <u>Note</u> )

Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.19 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human Histone H2A.X. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2025 / 11 / 26 Page 1 of 2




Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

### DATA IMAGES




#### GTX108272 IHC-P Image

Histone H2A.X antibody detects Histone H2A.X protein at nucleus in human esophagus cancer by immunohistochemical analysis.

Sample: Paraffin-embedded human esophagus cancer.

Histone H2A.X antibody (GTX108272) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




### GTX108272 IHC-P Image

Histone H2A.X antibody detects Histone H2A.X protein at nucleus in human cervical cancer by immunohistochemical analysis.

Sample: Paraffin-embedded human cervical cancer.

Histone H2A.X antibody (GTX108272) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




### GTX108272 IHC-P Image

Histone H2A.X antibody detects Histone H2A.X protein at nucleus on mouse prostate by immunohistochemical analysis.

Sample: Paraffin-embedded mouse prostate.

Histone H2A.X antibody (GTX108272) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



### GTX108272 IHC-P Image

Histone H2A.X antibody detects Histone H2A.X protein at nucleus on rat hind brain by immunohistochemical analysis.

Sample: Paraffin-embedded rat hind brain.

Histone H2A.X antibody (GTX108272) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our website.

Date 2025 / 11 / 26 Page 2 of 2