

N-RAS antibody

Cat. No. GTX108598

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human, Mouse, Rat

References (4) Package 100 μΙ, 25 μΙ

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	1:500-1:3000	
Not tested in other applications.		
Calculated MW	21 kDa. (<u>Note</u>)	
Product Note	Cross-reactivity may occur with other RAS proteins	

Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.41 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length human N-Ras Recombinant protein.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 12 / 03 Page 1 of 2




DATA IMAGES




GTX108598 WB Image

Various tissue extracts (50 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with N-RAS antibody (GTX108598) diluted at 1:1000. The HRP-conjugated anti-rabbit lgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX108598 WB Image

Various whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with N-RAS antibody (GTX108598) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX108598 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with N-RAS antibody (GTX108598) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 03 Page 2 of 2

€ 886-3-6208988 📻 886-3-6208989 🐷 infoasia@genetex.com