TKTL1 antibody [N1C1]

Cat. No. GTX109459

| Host | Rabbit |
|--------------|-------------------|
| Clonality | Polyclonal |
| lsotype | lgG |
| Applications | WB, ICC/IF, IHC-P |
| Reactivity | Human |
| Reactivity | Human |

References (4) Package 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|----------------------------------|----------------------|
| WB | 1:1000-1:10000 |
| ICC/IF | 1:100-1:1000 |
| IHC-P | 1:100-1:1000 |
| Not tested in other applications | |

Not tested in other applications.

| Calculated MW | 65 kDa. (<u>Note</u>) |
|---------------|---|
| Product Note | Based on internal testing result this antibody may cross react with human TKTL2 protein |

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS, 20% Glycerol |
| Preservative | 0.025% ProClin 300 |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 0.98 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Recombinant protein encompassing a sequence within the center region of human TKTL1. The exact sequence is proprietary. |
| Purification | Purified by antigen-affinity chromatography. |
| Conjugation | Unconjugated |



For full product information, images and publications, please visit our <u>website</u>.




Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX109459 ICC/IF Image

TKTL1 antibody [N1C1] detects TKTL1 protein at cytoplasm and nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: TKTL1 protein stained by TKTL1 antibody [N1C1] (GTX109459) diluted at 1:500. Red: Phalloidin, a cytoskeleton marker, diluted at 1:100. Scale bar = 10 μm.




GTX109459 IHC-P Image

Immunohistochemical analysis of paraffin-embedded SAS xenograft , using Transketolase-like protein 1(GTX109459) antibody at 1:100 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX109459 WB Image

Whole cell extract (30 µg) was separated by 7.5% SDS-PAGE, and the membrane was blotted with TKTL1 antibody [N1C1] (GTX109459) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX109459 WB Image

The data was published in the journal Aging Cell in 2014. PMID: 24757718



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 08 / 18 Page 2 of 2