

Factor X antibody

Cat. No. GTX109830

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 55 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Factor X. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX109830 ICC/IF Image

Immunofluorescence analysis of methanol-fixed HeLa, using Factor X(GTX109830) antibody at 1:200 dilution.



GTX109830 WB Image


Factor X antibody detects F10 protein by Western blot analysis.

A. 30 ug 293T whole cell lysate/extract

B. 30 ug whole cell lysate/extract of human F10-transfected 293T cells

10 % SDS-PAGE


Factor X antibody (GTX109830) dilution: 1:5000



GTX109830 IHC-P Image

Immunohistochemical analysis of paraffin-embedded A549 xenograft, using Factor X(GTX109830) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX109830 WB Image

HepG2 conditioned medium (30 μg) was separated by 10% SDS-PAGE, and the membrane was blotted with Factor X antibody (GTX109830) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](http://www.genetex.com).