

Fumarate hydratase antibody

Cat. No. GTX109877

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

References (3) Package 100 μΙ, 25 μΙ

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
Not tested in other applications.	

Calculated MW 55 kDa. (<u>Note</u>)

Properties	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.51 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Fumarate hydratase. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2025 / 12 / 07 Page 1 of 2




DATA IMAGES



GTX109877 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with Fumarate hydratase antibody (GTX109877) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX109877 ICC/IF Image

Fumarate hydratase antibody detects Fumarate hydratase protein at mitochondria by immunofluorescent analysis.

Sample: HeLa cells were fixed in ice-cold MeOH for 5 min.

Green: Fumarate hydratase stained by Fumarate hydratase antibody (GTX109877) diluted at 1:500. Blue: Fluoroshield with DAPI (GTX30920).

Scale bar= 10 µm.



GTX109877 WB Image


Fumarate hydratase antibody detects FH protein by western blot analysis.

A. 50 µg mouse liver lysate/extract

7.5% SDS-PAGE

Fumarate hydratase antibody (GTX109877) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX109877 IHC-P Image

Immunohistochemical analysis of paraffin-embedded NCIN87 xenograft, using fumarate hydratase (GTX109877) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 07 Page 2 of 2