

Tara antibody [C2C3-2], C-term

Cat. No. GTX110158

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Rat

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:10000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 261 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Tara. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

Date 2026 / 10 / 07 Page 1 of 2

DATA IMAGES

GTx110158 WB Image

Various whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Tara antibody [C2C3-2], C-term (GTx110158) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



GTx110158 IHC-P Image

Tara antibody [C2C3-2], C-term detects Tara protein at cytoplasm in human lung cancer by immunohistochemical analysis.


Sample: Paraffin-embedded human lung cancer.

Tara antibody [C2C3-2], C-term (GTx110158) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTx110158 WB Image

Whole cell extract (30 µg) was separated by 7.5% SDS-PAGE, and the membrane was blotted with Tara antibody [C2C3-2], C-term (GTx110158) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.


GTx110158 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using Tara(GTx110158) antibody at 1:200 dilution.



For full product information, images and publications, please visit our [website](https://www.genetex.com).