

## CD11c antibody [3.9]

Cat. No. GTX11029

<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Applications</b>	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, in vivo, Neutralizing/Inhibition
<b>Reactivity</b>	Human, Mouse, Rat, Cynomolgus monkey, Rhesus Monkey

References ( 8 )

Package

500 µg

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1-10 µg/mL
ICC/IF	Assay dependent
IHC-P	Assay dependent
IHC-Fr	Assay dependent
FCM	≤ 2 µg/10 <sup>6</sup> cells in a volume of 100 µL or 100µL of whole blood
IP	Assay dependent
in vivo	Assay dependent
Neutralizing/Inhibition	Assay dependent

Not tested in other applications.

**Calculated MW** 128 kDa. ( [Note](#) )

## Properties

<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 150 mM NaCl
<b>Preservative</b>	No preservatives
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	Batch dependent (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Rheumatoid synovial fluid cells and fibronectin purified human monocytes
<b>Purification</b>	Protein A/G purified



For full product information, images and publications, please visit our [website](#).

<b>Purity</b>	>95% by SDS-PAGE
<b>Endotoxin</b>	<0.5 EU/mg (determined by the LAL method)
<b>Conjugation</b>	Unconjugated
<b>Note</b>	<p>For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.</p> <p>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.</p>



For full product information, images and publications, please visit our [website](#).