

EZH2 antibody

Cat. No. GTX110384

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, ChIP assay
Reactivity	Human, Rat



APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
ChIP assay	Assay dependent
Not tested in other applications.	

Calculated MW 85 kDa. (<u>Note</u>)

PROPERTIES	
Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human EZH2. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2024 / 04 / 29 Page 1 of 2




DATA IMAGES



GTX110384 ChIP assay Image


Cross-linked ChIP was performed with HeLa chromatin extract and 5 µg of either control rabbit IgG or anti-EZH2 antibody. The precipitated DNA was detected by PCR with primer set targeting to CCND2.



GTX110384 ICC/IF Image

EZH2 antibody detects EZH2 protein at nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: EZH2 protein stained by EZH2 antibody (GTX110384) diluted at 1:500. Red: phalloidin, a cytoskeleton marker, diluted at 1:200. Blue: Hoechst 33342 staining.





GTX110384 WB Image

Sample (30 μg of whole cell lysate)

A: H1299

B: HCT116

C: MCF-7

7.5% SDS PAGE

GTX110384 diluted at 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.

For full product information, images and publications, please visit our website.

Date 2024 / 04 / 29 Page 2 of 2