

PYGM antibody [C1C3]

Cat. No. GTX110411

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse, Zebrafish

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 97 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.26 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human PYGM. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX110411 IHC-P Image

Immunohistochemical analysis of paraffin-embedded zebrafish tissue, using PYGM antibody [C1C3] (GTX110411) at 1:300 dilution.



GTX110411 WB Image

Sample (30 µg of whole cell lysate)

A: zebrafish eye

7.5% SDS PAGE

GTX110411 diluted at 1:1000



GTX110411 IHC-P Image

PYGM antibody [C1C3] detects PYGM protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse muscle.

PYGM stained by PYGM antibody [C1C3] (GTX110411) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTX110411 WB Image

Whole cell extract (30 µg) was separated by 7.5% SDS-PAGE, and the membrane was blotted with PYGM antibody [C1C3] (GTX110411) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).