

CDK8 antibody [C3], C-term

Cat. No. GTX110495

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

References (3)
Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 53 kDa. (Note)


Properties	
Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human CDK8. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 16 Page 1 of 2

DATA IMAGES




GTX110495 IHC-P Image

CDK8 antibody [C3], C-term detects CDK8 protein at cytoplasm and nucleus in mouse testis by immunohistochemical analysis.

Sample: Paraffin-embedded mouse testis.

CDK8 antibody [C3], C-term (GTX110495) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX110495 WB Image

CDK8 antibody detects CDK8 protein by western blot analysis. Various whole cell extracts ($50 \mu g$) were separated by 10% SDS-PAGE, and the membrane was blotted with CDK8 antibody (GTX110495) diluted at a dilution of 1:3000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 16 Page 2 of 2