

Destrin antibody

Cat. No. GTX11072

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat, Dog

References (2)

Package

50 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1500
ICC/IF	1:100
IHC-P	Assay dependent

Not tested in other applications.

Calculated MW 19 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA
Preservative	15mM Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	~0.5-1.0 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	synthetic peptide corresponding to amino acids 153-165 of human destriDF.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX11072 IHC-P Image

IHC-P analysis of rat stomach tissue using GTX11072 Destrin antibody at 0.02 µg/mL.


GTX11072 IHC-P Image

IHC-P analysis of mouse heart tissue using GTX11072 Destrin antibody at 0.02 µg/mL.


GTX11072 IHC-P Image

IHC-P analysis of mouse brain tissue using GTX11072 Destrin antibody at 0.02 µg/mL.



For full product information, images and publications, please visit our [website](#).