

CaMKII delta antibody

Cat. No. GTX111401

| Host | Rabbit |
|--------------|--------------------------------------|
| Clonality | Polyclonal |
| Isotype | IgG |
| Applications | WB, ICC/IF, IHC-P, IP |
| Reactivity | Human, Mouse, Rat, Zebrafish, Rabbit |



Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:50000 |
| ICC/IF | 1:100-1:1000 |
| IHC-P | 1:100-1:1000 |
| IP | 1:100-1:500 |

Not tested in other applications.

Calculated MW 56 kDa. (Note)

Product Note Cross-reactivity may occur with other CaM kinase II subunits.

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS, 20% Glycerol |
| Preservative | 0.025% ProClin 300 |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 0.66 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Recombinant protein encompassing a sequence within the center region of human CaMKII delta. The exact sequence is proprietary. |
| Purification | Purified by antigen-affinity chromatography. |
| Conjugation | Unconjugated |
| | |



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 13 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX111401 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (50 μ g) were separated by 10% SDS-PAGE, and the membrane was blotted with CaMKII delta antibody (GTX111401) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX111401 IHC-P Image

Immunohistochemical analysis of paraffin-embedded zebrafish tissue, using CaMKII delta antibody (GTX111401) at 1:300 dilution.



GTX111401 WB Image

Zebrafish tissue extract (30 μ g) was separated by 10% SDS-PAGE, and the membrane was blotted with CaMKII delta antibody (GTX111401) diluted at 1:40000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 06 / 13 Page 2 of 2