

CKMT2 antibody [N2C3]

Cat. No. GTX111620

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 48 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human CKMT2. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).


DATA IMAGES



GTx111620 IHC-P Image

Immunohistochemical analysis of paraffin-embedded BT483 xenograft, using sMtCK(GTx111620) antibody at 1:500 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTx111620 WB Image

Sample (30 ug of whole cell lysate)

A: 293T

10% SDS PAGE

GTx111620 diluted at 1:1000



GTx111620 IHC-P Image

CKMT2 antibody [N2C3] detects CKMT2 protein at mitochondria in mouse duodenum by immunohistochemical analysis.

Sample: Paraffin-embedded mouse duodenum.

CKMT2 antibody [N2C3] (GTx111620) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](https://www.genetex.com).