

TBCK antibody [C1C2], Internal

Cat. No. GTX111742

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 101 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.41 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human TBCK. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTx111742 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using TBCK(GTx111742) antibody at 1:200 dilution.


GTx111742 WB Image

Sample (30 µg of whole cell lysate)


A: HepG2 (GTx27900)

B: Molt-4 (GTx27912)

7.5% SDS PAGE


GTx111742 diluted at 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.


GTx111742 IHC-P Image

Immunohistochemical analysis of paraaffin-embedded MCF7 xenograft, using TBCK(GTx111742) antibody at 1:100 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


GTx111742 WB Image

Sample (30 µg of whole cell lysate)

A: NIH-3T3

B: JC

C: BCL-1

7.5% SDS PAGE

GTx111742 diluted at 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTx213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](http://www.genetex.com).