

# FLIP antibody [N1C1]

## Cat. No. GTX113047

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, IHC-P
Reactivity	Human

Reference (1) Package 100 μΙ, 25 μΙ

## APPLICATION

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

**Calculated MW** 55 kDa. ( Note )

PROPERTIES	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human FLIP. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our website.

Date 2024 / 05 / 03 Page 1 of 2




## DATA IMAGES



#### GTX113047 WB Image

Whole cell extract (30  $\mu$ g) was separated by 10% SDS-PAGE, and the membranes were blotted with FLIP antibody [N1C1] (GTX113047) diluted at 1:500 and competitor's antibody (sc-8347) diluted at 1:200. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody. \*The competitor is not affiliated with GeneTex and does not endorse this product.



#### GTX113047 IHC-P Image

Immunohistochemical analysis of paraffin-embedded human colon carcinoma, using FLIP(GTX113047) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 03 Page 2 of 2