

Fibrinogen gamma antibody

Cat. No. GTX113436

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 52 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.39 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Fibrinogen gamma. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX113436 IHC-P Image

Fibrinogen gamma antibody detects Fibrinogen gamma protein at cytosol on human colon carcinoma by immunohistochemical analysis.


Sample: Fibrinogen gamma antibody (GTX113436) dilution: 1:250.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min




GTX113436 ICC/IF Image

Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using Fibrinogen gamma (GTX113436) antibody at 1:200 dilution.



GTX113436 WB Image

Fibrinogen gamma antibody detects Fibrinogen gamma protein by Western blot analysis. Human plasma and HepG2 conditioned medium were separated by 10% SDS-PAGE, and the membrane was blotted with Fibrinogen gamma antibody (GTX113436) diluted at 1:2000.



GTX113436 WB Image

Human plasma (30 µg) was separated by 10% SDS-PAGE, and the membrane was blotted with Fibrinogen gamma antibody (GTX113436) diluted at 1:2000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).