FHL2 antibody

Cat. No. GTX114072

Host	Rabbit
Clonality	Polyclonal
lsotype	lgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

References (1) Package 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000
Not tested in other applications.	

Calculated MW

32 kDa. (<u>Note</u>)


Properties		
Form	Liquid	
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol	
Preservative	0.01% Thimerosal	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Recombinant protein encompassing a sequence within the center region of human FHL2. The exact sequence is proprietary.	
Purification	Purified by antigen-affinity chromatography.	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 13 Page 1 of 2


DATA IMAGES



GTX114072 IHC-P Image


Immunohistochemical analysis of paraffin-embedded SAS xenograft, using FHL2(GTX114072) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX114072 WB Image

Whole cell extract (30 μ g) was separated by 12% SDS-PAGE, and the membrane was blotted with FHL2 antibody (GTX114072) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX114072 WB Image

Mouse tissue extract (50 μ g) was separated by 12% SDS-PAGE, and the membrane was blotted with FHL2 antibody (GTX114072) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 13 Page 2 of 2