

Destrin antibody

Cat. No. GTX114170

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human, Mouse

References (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000

Not tested in other applications.

Calculated MW 19 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.39 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human Destrin. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES

GTX114170 WB Image

Sample (30 ug of whole cell lysate)
A: NIH-3T3
12% SDS PAGE
GTX114170 diluted at 1:2000


GTX114170 WB Image

Destrin antibody detects Destrin protein by Western blot analysis.
A. 30 μ g A549 whole cell lysate/extract
B. 30 μ g H1299 whole cell lysate/extract
C. 30 μ g HCT116 whole cell lysate/extract
12 % SDS-PAGE
Destrin antibody (GTX114170) dilution: 1:500


GTX114170 WB Image

Destrin antibody detects Destrin protein by Western blot analysis.
A. 30 μ g 293T whole cell lysate/extract
B. 30 μ g A431 whole cell lysate/extract
C. 30 μ g HeLa whole cell lysate/extract
D. 30 μ g HepG2 whole cell lysate/extract
12 % SDS-PAGE
Destrin antibody (GTX114170) dilution: 1:500



For full product information, images and publications, please visit our [website](https://www.genetex.com).