

# PDIA6 antibody

**Cat. No. GTX11432**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Application</b>	WB, ICC/IF, FACS
<b>Reactivity</b>	Human, Pig

**Package**

50 µl

## APPLICATION

### Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:250
ICC/IF	1:250
FACS	1:20

Not tested in other applications.

**Calculated MW** 48 kDa. ( [Note](#) )

## PROPERTIES

<b>Form</b>	Liquid
<b>Buffer</b>	Serum
<b>Preservative</b>	0.01% Sodium azide
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Immunogen</b>	Synthetic peptide corresponding to residues D(476) I D L S D V E L D D L G K D E L(492) of human P5.
<b>Purification</b>	Unpurified
<b>Conjugation</b>	Unconjugated


### Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).


**DATA IMAGES**

**GTX11432 WB Image**

WB analysis of membrane enriched extracts (30 µg lysate) of A549 (Lane 1), A-431 (Lane 2), HeLa (Lane 3), MCF7 (Lane 4), K-562 (Lane 5), Caco-2 (Lane 6), Hep G2 (Lane 7) and COLO 205 (Lane 8) using GTX11432 PDIA6 antibody.

Dilution : 1:250


**GTX11432 WB Image**

WB analysis of HT-1080 cell lysate using GTX11432 PDIA6 antibody.


**GTX11432 ICC/IF Image**

ICC/IF analysis of A549 cells using GTX11432 PDIA6 antibody. Panel e is a no primary antibody control.

Green : Primary antibody

Blue : Nuclei

Red : Actin

Fixation : 4% paraformaldehyde

Permeabilization : 0.1% Triton X-100 for 10 minutes

Dilution : 1:250 dilution in 0.1% BSA and incubated for 3 hours at room temperature



For full product information, images and publications, please visit our [website](https://www.genetex.com).