

RNF40 antibody [N1N2], N-term

Cat. No. GTX115295

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP
Reactivity	Human

References (1) Package $100 \mu l$, $25 \mu l$

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500

Not tested in other applications.

Calculated MW 114 kDa. (<u>Note</u>)

Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human RNF40. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2025 / 05 / 29 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX115295 WB Image


Sample (30 ug of whole cell lysate) A: IMR32 7.5% SDS PAGE GTX115295 diluted at 1:1000



GTX115295 IHC-P Image

Immunohistochemical analysis of paraffin-embedded human colon carcinoma, using RNF40(GTX115295) antibody at 1:250 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX115295 WB Image

RNF40 antibody detects RNF40 protein by western blot analysis. HeLa whole cell extracts and nuclear extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with RNF40 antibody (GTX115295) diluted by 1:1000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 05 / 29 Page 2 of 2

€ 886-3-6208988 💼 886-3-6208989 🖂 infoasia@genetex.com