

APC antibody [C3], C-term

Cat. No. GTX116009

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, IHC
Reactivity	Human, Mouse, Rat

References (3)

★★★★★ Review (1)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000
IHC	Assay dependent

Not tested in other applications.

Calculated MW 312 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS
Preservative	No preservatives
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence of human APC. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES


**GTX116009 IHC-P Image**

APC antibody [C3], C-term detects APC protein at cytoplasm by immunohistochemical analysis (Autostainer Formulated).

Sample: Paraffin-embedded human colon cancer.

APC stained by APC antibody [C3], C-term (GTX116009) diluted at 1:500.

Antigen Retrieval: Citrate buffer, 20 min


**GTX116009 IHC-P Image**

APC antibody [C3], C-term detects APC protein at cytoplasm in rat colon by immunohistochemical analysis.

Sample: Paraffin-embedded rat colon.

APC antibody [C3], C-term (GTX116009) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min


**GTX116009 IHC-P Image**

APC antibody [C3], C-term detects APC protein by immunohistochemical analysis.

Sample: Paraffin-embedded rat colon.

APC stained by APC antibody [C3], C-term (GTX116009) diluted at 1:200.

Antigen Retrieval: Tris-EDTA buffer, pH 9.0, 15 min

**GTX116009 WB Image**

Various whole cell extracts (50 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with APC antibody [C3], C-term (GTX116009) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident femto Western HRP Substrate.



For full product information, images and publications, please visit our [website](#).