

DDAH2 antibody [N2C3]

Cat. No. GTX116339

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 30 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human DDAH2. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX116339 WB Image

Sample (30 ug of whole cell lysate)
A: MCF-7
12% SDS PAGE
GTX116339 diluted at 1:1000



GTX116339 ICC/IF Image

DDAH2 antibody [N2C3] detects DDAH2 protein at mitochondria by immunofluorescent analysis.
Sample: SKNSH cells were fixed in 2% paraformaldehyde/culture medium at 37°C for 15 min.
Green: DDAH2 protein stained by DDAH2 antibody [N2C3] (GTX116339) diluted at 1:500.
Blue: Hoechst 33342 staining.
Scale bar = 10 μm.



GTX116339 IHC-P Image

DDAH2 antibody [N2C3] detects DDAH2 protein at mitochondria on human colon carcinoma by immunohistochemical analysis.
Sample: Paraffin-embedded colon carcinoma.
DDAH2 antibody [N2C3] (GTX116339) dilution: 1:500.
Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](#).