

URO4 antibody [S27]

Cat. No. GTX11770

Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Applications	ICC/IF, IHC-Fr, FCM
Reactivity	Human

References (1)

Package

500 µl

Applications

Application Note

FACS: Use at a dilution of 1/1 - 1/10. ICC: Use at a dilution of 1/40. Fix for 10 minutes in 4°C acetone. IHC-Fr: Use at a dilution of 1/40. Fix for 10 minutes in 4°C acetone. Optimal dilutions/concentrations should be determined by the end user.

Product Note This antibody detects a glycoprotein of 120kDa.

Properties

Form	Liquid
Buffer	PBS, 1% BSA
Preservative	0.1% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Immunogen	Tissue / cell preparation (Human SK-RC-7 Renal cancer cell line.).
Purification	Tissue culture supernatant
Conjugation	Unconjugated

Note For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).