

TRP1 antibody [Ta99]

Cat. No. GTX11780

Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Applications	ICC/IF, IHC-Fr
Reactivity	Human

References (1)

Package

500 µl

Applications

Application Note

IHC-Fr: Use at an assay dependent dilution. For optimal staining, the primary antibody should be incubated 20-60 minutes at roomtemperature. Optimal dilutions/concentrations should be determined by the end user.

Calculated MW 61 kDa. ([Note](#))

Product Note This monoclonal antibody detects a differentiation-related, pigmentation-associated glycoprotein (gp75), expressed by melanoma cells, nomal melanocytes and nevi. This antibody does not react with carcinomas and sarcomas.

Properties

Form	Liquid
Buffer	Preservative: 0.1% Sodium Azide; Constituents: 1% BSA, 0.01M PBS
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	Batch dependent (Please refer to the vial label for the specific concentration.)
Immunogen	SK-MEL-23 Melanoma cell line
Purification	Ammonium sulfate precipitation from tissue culture supernatant.
Conjugation	Unconjugated
Note	<p>For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.</p> <p>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.</p>



For full product information, images and publications, please visit our [website](#).