

STAT3 (phospho Tyr705) antibody

Cat. No. GTX118000

Host	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Applications	WB, ICC/IF, IHC-P, FCM	
Reactivity	Human, Mouse, Zebrafish	



Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

WB 1	1:500-1:3000
ICC/IF 1	1:100-1:1000
IHC-P A	Assay dependent
FCM A	Assay dependent

Not tested in other applications.

Calculated MW 88 kDa. (<u>Note</u>)

Properties		
Form	Liquid	
Buffer	PBS, 1% BSA, 20% Glycerol	
Preservative	0.025% ProClin 300	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Carrier-protein conjugated synthetic peptide surrounding phospho Tyr705 of human STAT3. The exact sequence is proprietary.	
Purification	Purified by antigen-affinity chromatography.	
Conjugation	Unconjugated	



For full product information, images and publications, please visit our website.

Date 2025 / 11 / 02 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX118000 WB Image

Untreated (–) and treated (+) A431 whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with STAT3 (phospho Tyr705) antibody (GTX118000) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX118000 ICC/IF Image

Confocal immunofluorescence analysis (Olympus FV10i) of paraformaldehyde-fixed A431, using STAT3 (Phospho Y705)(GTX118000) antibody (Green) at 1:500 dilution. Alpha-tubulin filaments were labeled with GTX11304 (Red) at 1:2000.



GTX118000 WB Image

Untreated (–) and treated (+) HeLa whole cell extracts (30 μ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with STAT3 (phospho Tyr705) antibody (GTX118000) diluted at 1:1000. The HRP-conjugated anti-rabbit lgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 02 Page 2 of 2