

# FLI1 antibody

**Cat. No. GTX11851**

|                     |              |
|---------------------|--------------|
| <b>Host</b>         | Sheep        |
| <b>Clonality</b>    | Polyclonal   |
| <b>Isotype</b>      | IgG          |
| <b>Applications</b> | WB           |
| <b>Reactivity</b>   | Human, Mouse |

**Package**  
125 µg

## Applications

### Application Note

WB: Use at a dilution of 1/500. This concentration is recommended using whole extracts from mouse erythroleukemia cell lines (CB7 and DP16-1) or a human T cell line (Jurkat). Detects a band of approximately 52 kDa. Optimal dilutions/concentrations should be determined by the end user.

**Calculated MW** 51 kDa. ( [Note](#) )

## Properties

|                      |  |
|----------------------|--|
| <b>Form</b>          | Liquid   |
| <b>Buffer</b>        | PBS  |
| <b>Preservative</b>  | 0.08% Sodium azide   |
| <b>Storage</b>       | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.   |
| <b>Concentration</b> | 1 mg/ml (Please refer to the vial label for the specific concentration.)   |
| <b>Immunogen</b>     | Fusion protein, corresponding to the N-terminal and amino acids 1-239 of Mouse FLI1.   |
| <b>Purification</b>  | IgG fraction<br>Whole antiserum is salt-fractionated to provide primarily the immunoglobulin fraction of antiserum.  |
| <b>Conjugation</b>   | Unconjugated   |
| <b>Note</b>          | For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.<br><br>Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product. |



For full product information, images and publications, please visit our [website](#).