

UPRT antibody

Cat. No. GTX120370

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

Package $100~\mu l, 25~\mu l$

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 34 kDa. (<u>Note</u>)


Properties	
Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human UPRT. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 30 Page 1 of 2


DATA IMAGES



GTX120370 IHC-P Image


Immunohistochemical analysis of paraffin-embedded human hepatoma, using UPRT(GTX120370) antibody at 1:500 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



GTX120370 WB Image

Sample (30 ug of whole cell lysate) A: A549 12% SDS PAGE GTX120370 diluted at 1:500



GTX120370 IHC-P Image

UPRT antibody detects UPRT protein at cytosol and nucleus on mouse liver by immunohistochemical analysis.

Sample: Paraffin-embedded mouse liver.

UPRT antibody (GTX120370) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 30 Page 2 of 2