

ADC antibody [N2C2], Internal

Cat. No. GTX120705

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Human

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:10000

Not tested in other applications.

Calculated MW 50 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.02 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of human ADC. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX120705 WB Image

ADC antibody detects ADC protein by western blot analysis.

A. 30 µg SK-N-SH whole cell lysate/extract

10 % SDS-PAGE

ADC antibody (GTX120705) dilution: 1:5000



GTX120705 WB Image

ADC antibody detects ADC protein by western blot analysis.

A. 30 µg human normal cerebellar lysate/extract

10 % SDS-PAGE

ADC antibody (GTX120705) dilution: 1:1500



For full product information, images and publications, please visit our [website](https://www.genetex.com).