

# IKK beta antibody [10AG2]

# Cat. No. GTX12139

Host	Mouse
Clonality	Monoclonal
Isotype	lgG1
Applications	WB, ICC/IF, IHC-P, FCM, IP, MS
Reactivity	Human, Mouse

Package 50 μg

# **Applications**

# **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	2 - 4 μg/ml
ICC/IF	Assay dependent
IHC-P	1:10 - 1:500
FCM	$0.25 - 0.5 \mu g  /  10^6  cells$
IP	1:10 - 1:500
MS	Assay dependent

Not tested in other applications.

**Calculated MW** 87 kDa. ( Note )

Properties	
Form	Liquid
Buffer	PBS
Preservative	0.05% Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	A full-length human IKKb recombinant protein was used to immunize mice (NP_001547).
Purification	Protein G purified
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 27 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


### DATA IMAGES



## GTX12139 FCM Image


FACS analysis of HEK293 cells using GTX12139 IKK beta antibody [10AG2].

Red: primary antibody Green: isotype control Antibody amount: 0.1µg



## GTX12139 WB Image

WB analysis of (A) Daudi, B) HeLa, and C) NIH3T3 cell lysate using GTX12139 IKK beta antibody [10AG2]. Dilution:  $2 \mu g/ml$ 



## GTX12139 FCM Image

FACS (Intracellular staining) analysis of HeLa cells using GTX12139 IKK beta antibody [10AG2].

Blue: Primary antibody Orange: isotype control Fixation: 4% PFA

Permibilization: 0.1% saponin Antibody amount: 1μg/100 μl buffer



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 27 Page 2 of 2