

Histone H3K9ac (Acetyl Lys9) antibody

Cat. No. GTX12178

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat, Drosophila, Bovine, Chicken, <i>Caenorhabditis elegans</i> , Frog

References (2)

Package

50 μ l

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1,000
ICC/IF	1:1,000
IHC-P	Assay dependent

Note : 20-40 μ g using RIPA extracts (0.45 mg) of rat PC12 cells

Not tested in other applications.

Properties	
Form	Liquid
Buffer	PBS
Preservative	15mM Sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	2-2.5 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	synthetic acetylated [Ac-Lys9] histone H3 peptide (amino acids 7-20) corresponding to the N-terminus of human histone H3. The sequence is identical in many species including mouse, rat, bovine, chicken, frog, Drosophila, and <i>C. elegans</i> , and is highly conserved (single amino acid substitution) in <i>Tetrahymena</i> histone H3.
Purification	Purified by affinity chromatography
Conjugation	Unconjugated




For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 1 of 2


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES**GTX12178 IHC-P Image**


IHC-P analysis of mouse testis tissue using GTX12178 Histone H3K9ac (acetyl Lys9) antibody at 0.1 µg/mL.

**GTX12178 IHC-P Image**

IHC-P analysis of rat lung tissue using GTX12178 Histone H3K9ac (acetyl Lys9) antibody at 0.1 µg/mL.

**GTX12178 IHC-P Image**

IHC-P analysis of mouse liver tissue using GTX12178 Histone H3K9ac (acetyl Lys9) antibody at 0.1 µg/mL.

**GTX12178 WB Image**

WB analysis of NIH3T3 cells using GTX12178 Histone H3K9ac (acetyl Lys9) antibody.

Lane 1 : Anti Histone H3K9ac (acetyl Lys9) antibody

Lane 2 : Anti Histone H3K9ac (acetyl Lys9) + histone H3 (acetylated Lys9) peptide

Lane 3 : Anti Histone H3K9ac (acetyl Lys9) + histone H3 peptide(non-acetylated)



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 2 of 2