Histone H3 antibody

Cat. No. GTX122150

Host	Rabbit
Clonality	Polyclonal
lsotype	lgG
Application	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Drosophila



APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:10000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

Not tested in other applications.


PROPERTIES	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of human Histone H3. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 18 Page 1 of 2

DATA IMAGES




GTX122150 WB Image


Whole cell extract (30 μ g) was separated by 15% SDS-PAGE, and the membrane was blotted with Histone H3 antibody (GTX122150) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX122150 ICC/IF Image

HIST1H3A antibody detects HIST1H3A protein at chromosome by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde/PBS for 15 min. Green: HIST1H3A protein stained by HIST1H3A antibody (GTX122150) diluted at 1:500. Blue: Hoechst 33342 staining.







GTX122150 IHC-P Image

Histone H3.1 antibody detects Histone H3.1 protein at nucleus on mouse intestine by immunohistochemical analysis. Sample: Paraffin-embedded mouse intestine.

Histone H3.1 antibody (GTX122150) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min

GTX122150 IHC-P Image

Histone H3.1 antibody detects Histone H3.1 protein at nucleus on mouse intestine by immunohistochemical analysis. Sample: Paraffin-embedded mouse intestine. Histone H3.1 antibody (GTX122150) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min

GTX122150 ICC/IF Image

Histone H3 antibody detects Histone H3 protein at nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: Histone H3 protein stained by Histone H3 antibody (GTX122150) diluted at 1:500. Red: phalloidin, a cytoskeleton marker, diluted at 1:50. Scale bar = 10 µm.



For full product information, images and publications, please visit our <u>website</u>.