

## Histone H4K5K8K12K16ac (acetyl Lys5/Lys8/Lys12/Lys16) antibody

**Cat. No. GTX122653**

<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	WB, ICC/IF, IHC-P, Dot
<b>Reactivity</b>	Human, Mouse, Rat

References ( 5 )

★★★★☆ Review ( 1 )

Package

100 µl, 25 µl

## Applications

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:5000-1:20000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
Dot	Assay dependent

Not tested in other applications.

## Properties


<b>Form</b>	Liquid
<b>Buffer</b>	PBS, 1% BSA, 20% Glycerol
<b>Preservative</b>	0.01% Thimerosal
<b>Storage</b>	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
<b>Concentration</b>	1 mg/ml (Please refer to the vial label for the specific concentration.)
<b>Immunogen</b>	Carrier-protein conjugated synthetic peptide surrounding acetyl Lys5/Lys8/Lys12/Lys16 of human Histone H4. The exact sequence is proprietary.
<b>Purification</b>	Purified by antigen-affinity chromatography.
<b>Conjugation</b>	Unconjugated

## Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

**DATA IMAGES**

**GTX122653 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded human breast cancer, using Histone H4K5K8K12K16ac (acetyl Lys5/Lys8/Lys12/Lys16)(GTX122653) antibody at 1:500 dilution. Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



**GTX122653 Dot Image**

Dotblot analysis of anti-Histone H4K5K8K12K16ac (acetyl Lys5/Lys8/Lys12/Lys16) antibody with peptide samples.

Peptide samples (0.1 µg) were spotted onto positively charged nylon membrane and blotted with Histone H4K5K8K12K16ac (acetyl Lys5/Lys8/Lys12/Lys16) antibody (GTX122653) at different dilution as indicated.

A: Peptide samples of Histone H4K5K8K12K16ac (acetyl Lys5/Lys8/Lys12/Lys16)

B: Peptide samples of Histone H4



**GTX122653 WB Image**

Sample (30 ug of whole cell lysate)

A: PC-12

15% SDS PAGE

GTX122653 diluted at 1:10000


**GTX122653 WB Image**

Sample (30 ug of whole cell lysate)

A: HeLa

15% SDS PAGE

GTX122653 diluted at 1:10000



For full product information, images and publications, please visit our [website](https://www.genetex.com).