

# TET2 antibody [N2-2], N-term

## Cat. No. GTX124205

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P, IP, ChIP assay
Reactivity	Human, Mouse

Reference (17) Package 100 μΙ, 25 μΙ

## APPLICATION

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500
ChIP assay	Assay dependent
Not tested in other applications.	

**Calculated MW** 224 kDa. ( Note )

PROPERTIES	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.55 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human TET2. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2024 / 04 / 28 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.


#### DATA IMAGES

Note



#### GTX124205 WB Image

Wild-type (WT) and TET2 knockout (KO) HeLa cell extracts (30  $\mu$ g) were separated by 5% SDS-PAGE, and the membrane was blotted with TET2 antibody [N2-2], N-term (GTX124205) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



#### GTX124205 WB Image

Various whole cell extracts (30  $\mu$ g) were separated by 5% SDS-PAGE, and the membrane was blotted with TET2 antibody [N2-2], N-term (GTX124205) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




### GTX124205 IHC-P Image

TET2 antibody [N2-2], N-term detects TET2 protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded human breast carcinoma.

TET2 stained by TET2 antibody [N2-2], N-term (GTX124205) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



#### GTX124205 ICC/IF Image

TET2 antibody [N2-2], N-term detects TET2 protein at nucleus by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: TET2 stained by TET2 antibody [N2-2], N-term (GTX124205) diluted at 1:500.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 04 / 28 Page 2 of 2