

Japanese encephalitis virus Envelope antibody

Cat. No. GTX125867

| Host | Rabbit |
|--------------|--|
| Clonality | Polyclonal |
| Isotype | lgG |
| Applications | WB, ICC/IF, IHC-P, IHC-P (cell pellet) |
| Reactivity | Japanese encephalitis virus |



Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

| Suggested dilution | Recommended dilution |
|---------------------|----------------------|
| WB | 1:5000-1:20000 |
| ICC/IF | 1:100-1:2000 |
| IHC-P | 1:100-1:1000 |
| IHC-P (cell pellet) | Assay dependent |

Not tested in other applications.

| Calculated | MW | 53 kDa. (| Note) |
|------------|----|-----------|--------|
| | | | |

Product Note

This antibody is specific for JEV Envelope protein, and it does not cross-react with Zika, DENV-1, DENV-2, DENV-3, and DENV-4 Envelope protein.

| Properties | |
|---------------|--|
| Form | Liquid |
| Buffer | PBS, 20% Glycerol |
| Preservative | 0.025% ProClin 300 |
| Storage | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| Concentration | 1.18 mg/ml (Please refer to the vial label for the specific concentration.) |
| Immunogen | Recombinant protein encompassing a sequence within the center region of Envelope protein (JEV). (Japanese Encephalitis Virus strain Jaoars982) The exact sequence is proprietary. |
| Purification | Purified by antigen-affinity chromatography. |
| Conjugation | Unconjugated |



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 14 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX125867 WB Image

Non-infected (–) and infected (+) BHK-21 whole cell extracts were separated by 12% SDS-PAGE, and the membrane was blotted with Japanese encephalitis virus Envelope antibody (GTX125867) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.




GTX125867 ICC/IF Image

Japanese encephalitis virus Envelope antibody detects Japanese encephalitis virus Envelope protein by immunofluorescent analysis.


Sample: Mock and transfected 293T cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: Japanese encephalitis virus Envelope stained by Japanese encephalitis virus Envelope antibody (GTX125867) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).



GTX125867 WB Image

Non-infected (–) and infected (+) BHK-21 whole cell extracts (5 μ g) were separated by 12% SDS-PAGE, and the membrane was blotted with Japanese encephalitis virus Envelope antibody (GTX125867) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX125867 ICC/IF Image

Envelope protein (Japanese encephalitis virus) antibody detects envelope protein (Japanese encephalitis virus) protein by immunofluorescent analysis.

Samples: BHK-21 cells mock (left) and infected with Japanese encephalitis viruswere fixed in MeOH. Green: envelope protein (Japanese encephalitis virus) protein stained by Envelope protein (Japanese encephalitis virus) antibody (GTX125867) diluted at 1:2000.

Blue: Hoechst 33342 staining.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 14 Page 2 of 2