# Japanese encephalitis virus NS3 antibody

# Cat. No. GTX125868

Host	Rabbit
Clonality	Polyclonal
lsotype	lgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr
Reactivity	Japanese encephalitis virus

References (33) Package 100 μl, 25 μl

## Applications

#### **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:5000-1:20000
ICC/IF	1:100-1:2000
IHC-P	Assay dependent
IHC-Fr	Assay dependent
Not tested in other applications.	

app

#### **Calculated MW**

69 kDa. (<u>Note</u>)

Properties	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.45 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the center region of NS3 (JEV). (Japanese Encephalitis Virus strain Jaoars982) The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated




For full product information, images and publications, please visit our website.

Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



#### GTX125868 WB Image

NS3 (Japanese encephalitis virus) antibody detects NS3 (Japanese encephalitis virus) protein by western blot analysis.

A. 10 µg BHK-21 whole cell extract


B. 10 µg whole cell extract of Japanese encephalitis virus infected BHK-21 cells

C. 10 µg whole cell extract of Dengue virus type 2 infected BHK-21 cells

10% SDS-PAGE

NS3 (Japanese encephalitis virus) antibody (GTX125868) dilution: 1:10000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




## GTX125868 ICC/IF Image

NS3 (Japanese encephalitis virus) antibody detects NS3 (Japanese encephalitis virus) protein by immunofluorescent analysis.

Samples: BHK-21 cells mock (left) and infected with Japanese encephalitis virus (right) were fixed in paraformaldehyde.

Green: NS3 (Japanese encephalitis virus) protein stained by NS3 (Japanese encephalitis virus) antibody (GTX125868) diluted at 1:2000.

Blue: Hoechst 33342 staining.



#### GTX125868 ICC/IF Image

NS3 (Japanese encephalitis virus) antibody detects NS3 (Japanese encephalitis virus) protein by immunofluorescent analysis.

Samples: BHK-21 cells mock (left) and infected with Japanese encephalitis viruswere fixed in MeOH. Green: NS3 (Japanese encephalitis virus) protein stained by NS3 (Japanese encephalitis virus) antibody (GTX125868) diluted at 1:2000. Blue: Hoechst 33342 staining.

For full product information, images and publications, please visit our <u>website</u>.