

Avian Influenza A virus H9N2 HA (Hemagglutinin) antibody

Cat. No. GTX127305

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB
Reactivity	Avian Influenza A virus

Package
100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000

Not tested in other applications.

Calculated MW 63 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.38 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of Influenza A virus H9N2 HA (Hemagglutinin) (A/turkey/Wisconsin/1/1966(H9N2)). The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX127305 WB Image

Influenza A Virus H9N2 Hemagglutinin (HA) antibody detects H9 (Influenza A) protein by Western blot analysis.

- A. MDCK cell lysates infected with 4 HAU/ml of A/duck/Alberta/35/1976-like, H1N1
- B. MDCK cell lysates infected with 4 HAU/ml of A/duck/Ukraine/1963-like, H3N8
- C. MDCK cell lysates infected with 4 HAU/ml of A/mallard/Miyagi/53/1976-like, H5N3
- D. MDCK cell lysates infected with 4 HAU/ml of A/duck/Taiwan/33/1993-like, H7N7
- E. MDCK cell lysates infected with 4 HAU/ml of A/turkey/Wisconsin/1/1966-like, H9N2
- F. MDCK cell lysates infected with 4 HAU/ml of A/PR/8/34, H1N1
- G. MDCK cell lysates infected with 4 HAU/ml of A/Taiwan/126/2009, pdmH1N1
- H. MDCK cell lysates infected with 4 HAU/ml of A/Taiwan/3446/2002, H3N2

12 % SDS-PAGE

Influenza A Virus H9N2 Hemagglutinin (HA) antibody (GTX127305) dilution: 1:1000



GTX127305 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Avian Influenza A virus H9N2 HA (Hemagglutinin) antibody (GTX127305) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).