

## 5-Hydroxymethylcytosine / 5-hmC antibody

Cat. No. GTX128454

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	Dot, MeDIP
Reactivity	Species independent

Package  
100 µl, 25 µl

## APPLICATION

## Application Note

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
Dot	Assay dependent
MeDIP	Assay dependent

Not tested in other applications.

## PROPERTIES

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	7.65 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	The immunogen used to generate this antibody corresponds to 5-HydroxymethylCytosine / 5-hmC
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


## Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.




For full product information, images and publications, please visit our [website](#).

**DATA IMAGES**

**GTX128454 MeDIP Image**

GTX128454 5-Hydroxymethylcytosine/ 5-hmC antibody in MeDIP experiment.

Human genomic DNA (150 ng) mixed with methylated cytosine standard kit (GTX400004) were subjected to MeDIP with 5-Hydroxymethylcytosine/ 5-hmC antibody (GTX128454) at dilution of 1:2000 and analyzed with semi-quantitative PCR. The immunoprecipitated DNA was plotted as % of input DNA.


**GTX128454 Dot Image**

GTX128454 5-hydroxymethylcytosine antibody Dot blot analysis of anti-5-hmC antibody with the synthetic DNA controls (GTX400004). DNA samples (0.5 to 50 ng as indicated) were spotted onto the positively charged Nylon membrane and blotted with 5-hmC antibody (GTX128454) at a dilution of 1:50000.

A: Unmethylated DNA

B: DNA containing 5-methylcytosine

C: DNA containing 5-hydroxymethylcytosine

D: DNA containing 5-formylcytosine

E: DNA containing 5-carboxylcytosine



For full product information, images and publications, please visit our [website](https://www.genetex.com).