

# mCherry antibody

## Cat. No. GTX128508

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-Fr, IHC-Wm, IP, PLA
Reactivity	Species independent



## APPLICATION

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:1000-1:10000
ICC/IF	1:100-1:1000
IHC-Fr	Assay dependent
IHC-Wm	1:100-1:500
IP	1:100-1:500
PLA	Assay dependent

Not tested in other applications.

Calculated MW 27 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.22 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Full length mCherry recombinant protein
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 03 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

#### DATA IMAGES




#### GTX128508 IHC-Wm Image

mCherry antibody labels mCherry-expressing cells on zebrafish embryos by whole mount immunohistochemical analysis.


Sample: 2 days-post-fertilization zebrafish embryo.

mCherry antibody labels mCherry-expressing cells (GTX128508) dilution: 1:100.




### GTX128508 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30  $\mu$ g) were separated by 7.5% SDS-PAGE, and the membrane was blotted with mCherry antibody (GTX128508) diluted at 1:5000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



#### GTX128508 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30  $\mu$ g) were separated by gradient gel% SDS-PAGE, and the membrane was blotted with mCherry antibody (GTX128508) diluted at 1:10000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



## GTX128508 ICC/IF Image

mCherry antibody detects mCherry protein at mCherry fusion protein by immunofluorescent analysis. Sample: mCherry fusion protein-transfected HeLa cells were fixed in 4% paraformaldehyde at RT for 30 min, then Triton X-100 for 1 min.

Green: mCherry protein stained by mCherry antibody (GTX128508) diluted at 1:500.

Red: mCherry is expressed in the transfected cell. Blue: Hoechst 33342 staining.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 03 Page 2 of 2