

DNMT3A antibody

Cat. No. GTX129125

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IP, EM
Reactivity	Human, Mouse, Rat



Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
IP	1:100-1:500
EM	Assay dependent
Not tested in other applications.	

Calculated MW 102 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.28 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human DNMT3A. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our website.

Date 2025 / 12 / 27 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note


Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX129125 WB Image

Non-transfected (–) and transfected (+) 293T whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with DNMT3A antibody (GTX129125) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.




GTX129125 WB Image

Wild-type (WT) and DNMT3A knockout (KO) 293T cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with DNMT3A antibody (GTX129125) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX129125 WB Image

Various whole cell extracts (30 μ g) were separated by 5% SDS-PAGE, and the membrane was blotted with DNMT3A antibody (GTX129125) diluted at 1:2000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX129125 IHC-P Image

DNMT3A antibody detects DNMT3A protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded mouse tissue.

DNMT3A stained by DNMT3A antibody (GTX129125) diluted at 1:500. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 27 Page 2 of 2