

CSDA antibody

Cat. No. GTX130052

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	WB, ICC/IF, IHC-P
Reactivity	Human, Mouse, Rat

Package
100 µl, 25 µl

APPLICATION

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	Assay dependent

Not tested in other applications.

Calculated MW	40 kDa. (Note)
Observed MW	40 48 kDa. The observed M.W. is based on the following publication. PMID: 26646932

PROPERTIES

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.99 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of human CSDA. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our [website](#).

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES




GTX130052 IHC-P Image

CSDA antibody detects CSDA protein at cytoplasm and nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded rat salivary gland.

CSDA stained by CSDA antibody (GTX130052) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX130052 IHC-P Image

CSDA antibody detects CSDA protein by immunohistochemical analysis.

Sample: Paraffin-embedded mouse tissues.

CSDA stained by CSDA antibody (GTX130052) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX130052 ICC/IF Image


CSDA antibody detects CSDA protein at cytoplasm by immunofluorescent analysis.

Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: CSDA stained by CSDA antibody (GTX130052) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).

Scale bar= 10µm.



GTX130052 WB Image

Rat tissue extract (50 µg) was separated by 10% SDS-PAGE, and the membrane was blotted with YBX3 antibody (GTX130052) diluted at 1:1000.

The observed M.W. is based on the publication: [PMID: 16603261](https://pubmed.ncbi.nlm.nih.gov/16603261/)



For full product information, images and publications, please visit our [website](https://www.genetex.com).