

# VASP antibody

# Cat. No. GTX132176

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P
Reactivity	Human



# Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000
Not tested in other applications	

**Calculated MW** 40 kDa. ( <u>Note</u> )


Properties	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	2.07 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the center region of human VASP. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our website.


Date 2025 / 11 / 07 Page 1 of 2

## DATA IMAGES



#### GTX132176 WB Image


Various whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with VASP antibody (GTX132176) diluted at1:1000.



#### GTX132176 ICC/IF Image

VASP antibody detects VASP protein at cytoplasm by immunofluorescent analysis. Sample: HeLa cells were fixed in 4% paraformaldehyde at RT for 10 min. Green: VASP protein stained by VASP antibody (GTX132176) diluted at 1:500. Blue: Hoechst 33342 staining.

Scale bar =  $10 \mu m$ .




#### GTX132176 IHC-P Image

VASP antibody detects VASP protein at cytoplasm in MDA-MB-231 xenograft by immunohistochemical analysis.

Sample: Paraffin-embedded MDA-MB-231 xenograft.

VASP antibody (GTX132176) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



### GTX132176 WB Image

Untreated (–) and treated (+) 293T whole cell extracts (30  $\mu$ g) were separated by 10% SDS-PAGE, and the membrane was blotted with VASP antibody (GTX132176) diluted at 1:2000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 11 / 07 Page 2 of 2