p53 (phospho Ser15) antibody

Cat. No. GTX132995

Host	Rabbit		References (
Clonality	Polyclonal		***
lsotype	IgG		<mark>Раскаде</mark> 100 µl, 25 µl
Applications	WB, ICC/IF		
Reactivity	Human		

References (1)
★ ★ 🚖 ★ Review (1)
<mark>Package</mark> 100 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	1:500-1:3000	
ICC/IF	1:100-1:1000	
Not tested in other applications.		

Calculated MW

44 kDa. (<u>Note</u>)


Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.23 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide surrounding phospho Ser15 of human p53. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 09 / 08 Page 1 of 2

DATA IMAGES




GTX132995 ICC/IF Image

p53 (phospho Ser15) antibody detects p53 (phospho Ser15) protein at nucleus by immunofluorescent analysis.


Sample: Mock and treated 293T cells were fixed in 4% paraformaldehyde at RT for 15 min. Green: p53 (phospho Ser15) protein stained by p53 (phospho Ser15) antibody (GTX132995) diluted at 1:2000. Blue: Hoechst 33342 staining.

Scale bar = 10 µm.



GTX132995 WB Image

Untreated (–) and treated (+) 293T whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with p53 (phospho Ser15) antibody (GTX132995) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX132995 WB Image

Untreated (–) and treated (+) 293T whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with p53 (phospho Ser15) antibody (GTX132995) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 09 / 08 Page 2 of 2