

# NeuN antibody

# Cat. No. GTX133127

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Application	ICC/IF, IHC-P, IHC-Fr
Reactivity	Mouse, Rat

Reference (3)
Package
100 µl, 25 µl

## APPLICATION

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	1:100-1:1000
IHC-Fr	1:100-1:1000
Not tested in other applications	

Not tested in other applications.

Calculated MW 41 kDa. (Note)

PROPERTIES	
Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.26 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the N-terminus region of mouse NeuN. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 02 Page 1 of 2




## DATA IMAGES



#### GTX133127 IHC-P Image

NeuN antibody detects NeuN protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded mouse hippocampus. NeuN stained by NeuN antibody (GTX133127) diluted at 1:1000. Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



### GTX133127 ICC/IF Image

NeuN antibody detects NeuN protein at nucleus by immunofluorescent analysis.

Sample: DIV9 rat E18 primary hippocampal neuron cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: NeuN stained by NeuN antibody (GTX133127) diluted at 1:250.

Red: Tau, a axon marker, stained by Tau antibody [GT287] (GTX634809) diluted at 1:500.




#### GTX133127 IHC-P Image

NeuN antibody detects NeuN protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded rat cerebellum.

NeuN stained by NeuN antibody (GTX133127) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




#### GTX133127 IHC-P Image

NeuN antibody detects NeuN protein at nucleus by immunohistochemical analysis. Sample: Paraffin-embedded rat hippocampus.

NeuN stained by NeuN antibody (GTX133127) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



### GTX133127 IHC-Fr Image

NeuN antibody detects NeuN protein at cytoplasm and nucleus by immunohistochemical analysis.

Sample: Frozen-sectioned mouse spinal cord.

Green: NeuN stained by NeuN antibody (GTX133127) diluted at 1:500.

Antigen Retrieval: 4% paraformaldehyde



For full product information, images and publications, please visit our <u>website</u>.

Date 2024 / 05 / 02 Page 2 of 2