

# Aquaporin 4 antibody

## Cat. No. GTX133151

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, ICC/IF, IHC-P, IHC-Fr, IHC (Free Floating)
Reactivity	Mouse, Rat



## Applications

## **Application Note**

\*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution	
WB	1:500-1:3000	
ICC/IF	1:100-1:1000	
IHC-P	1:100-1:1000	
IHC-Fr	1:100-1:1000	
IHC (Free Floating)	Assay dependent	
Not tested in other applications.		

Calculated MW 34 kDa. (Note)

Properties	
Form	Liquid
Buffer	PBS, 1% BSA, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.31 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the Intracellular domain of mouse Aquaporin 4. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 02 Page 1 of 2




For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

#### Note

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

### DATA IMAGES




#### GTX133151 IHC-P Image

Aquaporin 4 antibody detects Aquaporin 4 protein at cell membrane in mouse brain by immunohistochemical analysis.

Sample: Paraffin-embedded mouse brain.

Aquaporin 4 antibody (GTX133151) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



## GTX133151 ICC/IF Image

Aquaporin 4 antibody detects Aquaporin 4 protein at astrocyte cells by immunofluorescent analysis.

Sample: DIV9 rat E18 primary cortical neurons and glia cells were fixed in 4% paraformaldehyde at RT for 15 min.

Green: Aquaporin 4 protein stained by Aquaporin 4 antibody (GTX133151) diluted at 1:500. Red: beta Tubulin 3/ Tuj1, a neuron cell marker, stained by beta Tubulin 3/ Tuj1 antibody [GT886] (GTX631830) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).




## GTX133151 IHC-P Image

Aquaporin 4 antibody detects Aquaporin 4 protein at cell membrane by immunohistochemical analysis. Sample: Paraffin-embedded rat brain.

Aquaporin 4 stained by Aquaporin 4 antibody (GTX133151) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



#### GTX133151 IHC-Fr Image

Aquaporin 4 antibody detects Aquaporin 4 protein expression by immunohistochemical analysis. Sample: Frozen-sectioned adult mouse cerebellum.

Green: Aquaporin 4 protein stained by Aquaporin 4 antibody (GTX133151) diluted at 1:250.

Red: NeuN, stained by NeuN antibody [2Q158] (GTX30773) diluted at 1:500.

Blue: Fluoroshield with DAPI (GTX30920).



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 12 / 02 Page 2 of 2