EGFR (phospho Ser1046/Ser1047) antibody

Cat. No. GTX133433

ClonalityPolyclonalIsotypeIgGApplicationsWBReactivityHuman	Host	Rabbit
Applications WB	Clonality	Polyclonal
	lsotype	lgG
Reactivity Human	Applications	WB
	Reactivity	Human

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Optimal dilutions/concentrations should be determined by the researcher.		
Suggested dilution	Recommended dilution	
WB	1:500-1:3000	
Not tested in other applications.		
Calculated MW	134 kDa. (<u>Note</u>)	
Product Note	The full-length EGFR includes the 24 aa signal peptide that is cleaved during EGFR maturation. The Ser1046/Ser1047 of mature EGFR is equivalent to Ser1070/Ser1071 of uncleaved EGFR.	
Properties		
Form	Liquid	
Buffer	PBS, 1% BSA, 20% Glycerol	
Preservative	0.025% ProClin 300	
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.	
Concentration	0.31 mg/ml (Please refer to the vial label for the specific concentration.)	
Immunogen	Carrier-protein conjugated synthetic peptide surrounding phospho Ser1046/Ser1047 of human EGFR. The exact sequence is proprietary.	
Purification	Purified by antigen-affinity chromatography.	
Conjugation	Unconjugated	
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.	




For full product information, images and publications, please visit our website

publications, please visit our <u>website</u>.

🐛 886-3-6208988 💼 886-3-6208989 🔜 infoasia@genetex.com

Date 2025 / 07 / 17 Page 1 of 2

DATA IMAGES



GTX133433 WB Image

Untreated (–) and treated (+) A431 whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with EGFR (phospho Ser1046/Ser1047) antibody (GTX133433) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 07 / 17 Page 2 of 2