

NLRP3 antibody

Cat. No. GTX133569

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Mouse

References (4)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	Assay dependent

Not tested in other applications.

Calculated MW 118 kDa. ([Note](#))

Properties

Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1.57 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the N-terminus region of human NLRP3. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated

Note


For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

DATA IMAGES




GTX133569 IHC-P Image

NLRP3 antibody detects NLRP3 protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse spleen.


NLRP3 stained by NLRP3 antibody (GTX133569) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min




GTX133569 WB Image

Boiled and unboiled THP-1 whole cell and membrane extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with NLRP3 antibody (GTX133569) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody. (WCE: whole cell extract; ME: membrane extract)



GTX133569 WB Image

Untreated (-) and treated (+) THP-1 whole cell extract (50 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with NLRP3 antibody (GTX133569) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



GTX133569 WB Image

Untreated (-) and treated (+) Raw264.7 whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with NLRP3 antibody (GTX133569) diluted at 1:500. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](https://www.genetex.com).