

Doublecortin antibody

Cat. No. GTX134052

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P, IHC-Fr
Reactivity	Human, Mouse, Rat

References (6)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000
IHC-Fr	1:100-1:1000

Not tested in other applications.

Calculated MW 41 kDa. ([Note](#))

Properties


Form	Liquid
Buffer	PBS, 20% Glycerol
Preservative	0.025% ProClin 300
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.72 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Carrier-protein conjugated synthetic peptide encompassing a sequence within the C-terminus region of mouse Doublecortin. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated
For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.	
Note	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 1 of 2


DATA IMAGES



GTGX134052 WB Image


Various tissue extracts (50 µg) were separated by 10% SDS-PAGE, and the membranes were blotted with Doublecortin antibody (GTGX134052) diluted at 1:1000 and competitor's antibody (MO22113) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTGX213110-01) was used to detect the primary antibody.

*Competitor's antibody is not affiliated with GeneTex and does not endorse this product.




GTGX134052 WB Image

Various whole cell extracts (30 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with Doublecortin antibody (GTGX134052) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTGX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTGX134052 WB Image

Various tissue extracts (50 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with Doublecortin antibody (GTGX134052) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTGX213110-01) was used to detect the primary antibody.



GTGX134052 IHC-P Image

Doublecortin antibody detects Doublecortin protein at cytoplasm by immunohistochemical analysis.

Sample: Paraffin-embedded mouse E10.5 mouse embryo.

Doublecortin stained by Doublecortin antibody (GTGX134052) diluted at 1:1000.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).

Date 2026 / 02 / 01 Page 2 of 2